
No.



HHE UNAHED SHARES OF A

TO ALL TO WHOM THESE PRESENTS SHALL COME; Thio Agricultural Research Development Center

Telliereas, there has been presented to the

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED NOVEL VARIETY OF SEXUALLY REPRODUCED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED to be entitled to a certificate of plant variety protection under the ${
m LAW}.$

NOW, THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OR ASSIGNS OF THE SAID APPLI-CANT(S) FOR THE TERM OF Seventeen YEARS FROM THE DATE OF THIS GRANT, SUBJECT TO THE PAYMENT OF THE REQUIRED FEES AND PERIODIC REPLENISHMENT OF VIABLE BASIC SEED OF THE VARIETY IN A PUBLIC REPOSITORY AS PROVIDED BY LAW, THE RIGHT TO EX-CLUDE OTHERS FROM SELLING THE VARIETY, OR OFFERING IT FOR SALE, OR REPRODUCING IT. OR IMPORTING IT, OR EXPORTING IT, OR USING IT IN PRODUCING A HYBRID OR DIFFERENT ariety therefrom, to the extent provided by the Plant Variety Protection Act.THE UNITED STATES SEED OF THIS VARIETY (1) SHALL BE SOLD BY VARIETY NAME ONLY AS ASS OF CERTIFIED SEED AND (2) SHALL CONFORM TO THE NUMBER OF GENERATIONS OWNER OF THE RIGHTS. (84 STAT. 1542, AS AMENDED, 7 U.S.C. 2321 ET SEQ.)

WHEAT

'Ruler'

In Testimony Withereot, Shave hereunto set my hand and caused the seal of the Maut Variety Protection Office to be affixed at the City of Washington this 20th day of November in the year of our Lord one thousand nine hundred and seventy-four

and Bit

UNITED STATES DEPARTMENT OF AGRICULTURE
AGRICULTURAL MARKETING SERVICE
GRAIN DIVISION
HYATTSVILLE, MARYLAND 20782 FORM GR - 470 (12-15-72)

FORM APPROVED OMB NO. 40-R3712

APPLICATION FOR PLANT VARIETY PROTECTION CERTIFICATE

STRUCTIONS: See Reverse. VARIETY NAME OR TEMPORARY	2. KIND NAME		FOR OFFICIAL USE ONLY	
DESIGNATION	TI RAME ON TEM ONLY		PV NUMBER 7 4000 78	
GENUS AND SPECIES NAME	4. FAMILY NAME (B)	otanical)	FILING DATE	TIME
GENUS AND SPECIES NAME	10 1 11 11 11 11 11 11	-	13/18.74	3;00 F.M.
Triticum <u>aestivum</u> L.	Gramineae		FEE RECEIVED	BALANCE DUE
2034144111	5. DATE OF DETERMINATION		s 250,00	\$
	July 20, 1	1973	\$250.00	\$
·			\$250.00	\$
NAME OF APPLICANT(S)	7. ADDRESS (Street . Code)	and No. of R.F.D. No.,	City, State, and ZIP	8. TELEPHONE AREA CODE AND NUMBER
Ohio Agricultural Research				216 264 1221
Development Center	Wooster, Ohi	to 44691		216-264-1021
			•	
IF THE NAMED APPLICANT IS NOT A PE	RSON, FORM OF	10. STATE OF INCO	RPORATION	11. DATE OF INCOR-
ORGANIZATION: (Corporation, partnership,	association, etc.)			PORATION
Agriculture Experiment Stati	on	Ohio	·	
Name and mailing address of applic	ant representative	(s), if any, to serve	in this application a	nd receive all papers
Agronomy Department Ohio Agricultural Res. & Dev Mooster, Ohio 44691	. Center			
X 13A. Exhibit A, Origin and Bree	cription of the Vari	ety	ion)2 of the Flam Vo	artery Protection con-
X 130. Exhibit C, Objective Desc	ription of the Varie	ety		
X 13D. Exhibit D, Data Indicative	e of Novelty			
X 13E. Exhibit E, Statement of th	e Basis of Applica	nt's Ownership	·	
14A. Does the applicant(s) specify tha (See Section 83(a), (If "Yes," an	t seed of this varie	ty be sold by varie	ty name only as a cla	ass of certified seed?
148. Does the applicant(s) specify tha	t this variety be	14C. If "Yes," t	o 14B, how many gen	erations of production
limited as to number of generatio		beyond bree	eder seed?	
Timeter and it in amount in Parisana	XYES NO	o 🙀 FOUNDAT	ION REGISTERE	D CERTIFIE
The applicant declares that a viable ance of a certificate and will be replaced	sample of basic sec	ed of this variety w	vill be deposited upon th such regulations a	request before issus
	emoned periodica.	ced novel plant var	iety helieves that the	variety is distinct.
ance of a certificate and with both pro-	a anno allo abbandes	cea novel prant van	iety delieves that the	Dunius,
The undersigned applicant(s) of this uniform, and stable as required in Stable as required in Stable as required in Stable as required in Stable 2015.	ection 41 and is en	ntitled to protection	n under the provisions	s of Section 42 of the
The undersigned applicant(s) of thi uniform, and stable as required in S	ection 41 and is en	ntitled to protection	n under the provisions	s of Section 42 of the
The undersigned applicant(s) of thi uniform, and stable as required in S Plant Variety Protection Act.	ection 41 and is en	can jeopardize prot	tection and result in p	penalties.

INSTRUCTIONS

GENERAL: Send an original copy of the application, exhibits and \$250.00 fee to U.S. Dept. of Agriculture, Agricultural Marketing Service, Grain Division, 6525 Belcrest Road, Hyattsville, Maryland 20782. (See Section 180.175 of the regulations and rules of practice.) Retain one copy for your files. All items on the face of the form are self-explanatory unless noted below.

ITEM

- Insert the date the applicant determined that he had a new variety based on the definition in Section 41 (a) of the Act and decision is made to increase the seed.
- 13a First, give the genealogy, including public and commercial varieties, lines, or clones used, and the breeding method. Second, give the details of subsequent stages of selection and multiplication. Third, indicate the type and frequency of variants during reproduction and multiplication and state how these variants may be identified. Fourth, provide evidence on stability.
- 13b First, give any special characteristics of the seed and of the plant as it passes through the seedling stage, flowering stage and the fruiting stage. Second, describe the mature plant and compare it with a similar commercial variety grown under the same conditions, and indicate the differences.
- 13c A supplemental form will be furnished by the PVPO to describe in detail a variety for each kind of seed.
- 13d Provide complete data indicative of novelty. Seed and plant specimens or photographs of seed and plant comparisons clearly indicating novelty may be submitted. Seeds submitted may be sterile.
- 13e Indicate whether applicant is the actual breeder, the employer of the breeder, the owner through purchase or inheritance, etc.

Exhibit A

Origin and Breeding History of the Variety

- 1. TN 1542 (temporary designation) originated in Ohio from a hand-pollinated cross of a Purdue University line known as 'L494Al-8-5-5' by 'Lucas' using Lucas as the pollen parent. The cross was made in 1957 and was designated '757'. L494Al-8-5-5 was a line selected from the cross of 'Norin 10-3' by 'Purdue R4746A6'. Purdue R4746A6 originated from a complex cross of (Kawvale-Fultz-Hungarian-W38-Wabash-Fairfield-Trumbull³-Hope-Hussar) X (Trumbull-Fultz sel.-Minhardi-Wabash-Purplestraw-Chinese-Michigan Amber). Lucas was an Ohio-developed variety and the breeding history of Lucas may be found in Technical Bulletin 1278, U.S.D.A. entitled "Classification of Triticum Species and of Wheat Varieties Grown in the United States".
- 2. The line traces back to a single head selection made in the F₆ generation in 1963. The line was advanced to replicated yield trials in 1966 at Wooster, Ohio only, two locations in 1967, four locations in 1968, 5 locations in 1969, and at least 7 locations each succeeding year through 1973, all in Ohio. Further purification of the line was made by making a total of 208 head selections in 1967, 1970, and 1972. Breeders seed was comprised of the bulk of the progeny of these latter head selections. The line was first entered in the Uniform Eastern Soft Wheat Nursery in 1971 and continued in 1972 and 1973.
- 3. The line appears to be very homogeneous as observed in the field for the past 3 generations. This would logically be expected as the line was developed from a single head selected in F_6 with further selection in the F_{10} , F_{13} and F_{15} generations.
- 4. The line appears to be very stable and true breeding. However, some sterility occurs on TN-1542 under unusual environmental conditions which can lead to the occurrence of off-types the following generation due to outcrossing with adjacent varieties. The type of outcrosses varies depending on what variety is adjacent when these conditions occur. Most off-types can be expected to resemble TN 1542 as this line would be serving as the maternal parent. The number of off-types would be dependent on the proximity and maturity of adjacent varieties when sterility conditions occur, but have been observed to be minimal.

Exhibit B

Botanical Description of the Variety

RULER THE TWO TRY 1542 is a very short, midseason soft red winter wheat with a dark green foliage. The stem is stiff and erect. At maturity the head is only slightly nodding. At heading the flag leaf is semi-erect. Winterhardiness under Ohio conditions is excellent. Early fall growth is mostly prostrate to semi-erect. Tillering ability is very good. TN 1542 appears very vigorous under most growth conditions.

RULERANE TN 1542 is moderately tolerant to acid-soil conditions, very resistant to soil-borne spindle streak mosaic virus and moderately resistant to loose smut, but susceptible to most races of leaf rust, stem rust, and moderately susceptible to powdery mildew. It is also resistant to the Great Plains, A, C, and F races of Hessian fly.

RULFRATE
TN-1542 heads about 5 days later than Arthur, produces large, mid-dense, fusiform heads with yellow anthers and round-shoulder glumes of medium length which are awnleted. Tip awns are from 1 to 3 cm. in length.

FORM GR-470-6 (2-15-73)

UNITED STATES DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE GRAIN DIVISION HYATTSVILLE, MARYLAND 20782

EXHIBIT C (Wheat)

OBJECTIVE DESCRIPTION OF VARIETY WHEAT (TRITICUM SPP.)

INSTRUCTIONS: See Reverse. WHEAT (TRI)	
NAME OF APPLICANT(S)	FOR OFFICIAL USE ONLY
Ohio Agricultural Research & Development Condition of R.F.D. No., City, State, and ZIP Code)	enter PVPO NUMBER 7460078 146
•	VARIETY NAME OR TEMPORARY
Wooster, Ohio 44691	DESIGNATION
	Rulerato
Place the appropriate number that describes the varietal character Place a zero in first box (e.g. 0 8 9 or 0 9) when number i	of this variety in the boxes below. s either 99 or less or 9 or less.
1. KIND:	
1 1 = COMMON 2 = DURUM 3 = EMMER 4 = SPELT 5	= POLISH 6 = POULARD 7 = CLUB
2. TYPE: 2 1 = SPRING 2 = WINTER 3 = OTHER (Specify)	1 = SOFT 3 = OTHER (Specify) 1 2 = HARD
- Commence of the commence of	
2 1 = WHITE 2 = RED 3 = OTHER (Specify)	
3. SEASON - NUMBER OF DAYS FROM EMERGENCE TO:	
2 3 0 FIRST FLOWERING	2 3 5 LAST FLOWERING
4. MATURITY (50% Flowering):	
NO. OF DAYS EARLIER THAN	1 = ARTHUR 2 = SCOUT 3 = CHRIS
0 5 NO. OF DAYS LATER THAN	4 = LEMHI 5 = NUGAINES 6 = LEEDS
5. PLANT HEIGHT (From soil level to top of head):	
0 9 9 cm. HIGH	
0 4 CM. TALLER THAN	
CM. SHORTER THAN	4 = LEMHI 5 = NUGAINES 6 = LEEDS
6. PLANT COLOR AT BOOTING (See reverse):	7. ANTHER COLOR:
3 1 = YELLOW GREEN 2 = GREEN 3 = BLUE GREEN	1 = YELLOW 2 = PURPLE
8, STEM:	
1 Anthocyanin: 1 = ABSENT 2 = PRESENT	Waxy bloom: = ABSENT 2 = FRESENT
Hairiness of last internode of rachis: 1 = ABSENT 2 = PRESENT	1 Internodes: 1 = HOLLOW 2 = SOLID
0 4 NO. OF NODES (Originating from node above ground)	CM. INTERNODE LENGTH BETWEEN FLAG LEAF AND LEAF BELOW
9. AURICLES:	
Anthocyanin: 1 = ABSENT 2 = PRESENT	Hairiness: = ABSENT 2 = PRESENT
10. LEAF:	
Flag leaf at 1 = ERECT 2 = RECURVED booting stage: 3 = OTHER (Specify): Semi-erect	1 Flag leaf: 1 = NOT TWISTED 2 = TWISTED
2 Hairs of first leaf sheath: 1 = ABSENT 2 = PRESENT	1 Waxy bloom of flag leaf sheath: 1 = ABSENT 2 = PRESENT
1 2 MM. LEAF WIDTH (First leaf below flag leaf)	2 7 CM. LEAF LENGTH (First leaf below flag leaf):

Exhibit D

Data Indicative of Novelty

RULER of Novelty

TN-1542 is a soft red winter wheat which most closely resembles Logan and Lucas, however, it is about 10 cm shorter, 2 days earlier, has stiffer straw, is more susceptible to leaf rust, and most resistant to wheat spindle streak mosaic virus than Logan. TN 1542 is about 4 days earlier and 15 cm shorter than Lucas. than Lucas.

Office 31, Blaak P.O.B. 182

Telephone 010 - 142211 Telegrams Cebeco Telex No. 21398 P.C. account 5658

Bank: Centrale Rabobank

Head-Office Utrecht No. of account 30.00.00.065



CEBECO-HANDELSRAAD

Nationale coöperatieve aan- en verkoopvereniging voor land- en tuinbouw g.a. National agricultural co-operative wholesale society

· By Airmail

U.S. Department of Agriculture
Agricultural Marketing Service, Grain Division
National Agricultural Library
BELTSVILLE - Maryland 20705
U.S.A.

Dept.: Seed

dW/vG.

Extension:

ROTTERDAM,

278

August 31st, 1976.

Gentlemen:

We refer to your letter of July 30, 1976 from which we learned that you were kind enough to refer our request for 2 kg. seed sample of "Ruler" wheat to Dr. Joseph Craddock, Plant Genetics and Germplasm Institute, ARS, Beltsville, Md. 20705.

This will thank you for your kindness and acknowledge due receipt of the 2 kg. of "Ruler" winter wheat. We shall be glad if you inform Dr. Craddock of the sample having been received.

Very truly yours,

CEBECO-HANDELSRAAD

mod. 910.05

Exhibit D

Data Indicative of Novelty

*RULER WE
TN 1542 is a soft red winter wheat which most closely resembles Logan and Lucas, however, it is about 10 cm shorter, 2 days earlier, has stiffer straw, is more susceptible to leaf rust, and most resistant to wheat spindle streak mosaic virus than Logan. TN 1542 is about 4 days earlier and 15 cm shorter than Lucas.

Exhibit E

Statement of Applicant's Ownership

The Ohio Agricultural Research and Development Center, Wooster, Ohio is the sole, original and first developer and owner of $\frac{70.01542}{10.01542}$ variety of soft red winter wheat for which it solicits a certificate of protection.

FORM GR-470-6 (REVERSE)					
11. HEAD:	· · · · · · · · · · · · · · · · · · ·		0 2		
Density: 1 = LAX 2	= DENSE 3=(Mid-dense)	Shape: 1 = TAPERING 1 = OTHER (Sp.	2 = STRAP 3 = CLAVATE ecify)		
3 Awnedness: 1 = AWNLESS 2 = APICALLY AWNLETED 3 = AWNLETED 4 = AWNED					
7 Color at maturity: 5 = BF	HITE 2 = YELLOW 3 = PINK 4 = R ROWN 6 = BLACK 7 = OTHER	Specify): white to yel	1ow		
1 1 CM. LENGTH		1 1 MM. WIDTH			
12. GLUMES AT MATURITY:		 -			
2 Length: 1 = SHORT (CA 3 = LONG (CA.		Width: 1 ≈ NARROW (C 3 = WIDE (CA.			
Shoulder 1 = WANTING shape: 4 = SQUARE		Beak: 1 = OBTUSE	2 = ACUTE 3 = ACUMINATE		
13. COLEOPTILE COLOR:		14. SEEDLING ANTHOCYAN	M;		
1 I = WHITE 2 = RED	3 = PURPLE	2 1 = ABSENT 2 =	PRESENT		
15. JUVENILE PLANT GROW	TH HABIT:				
2 1 = PROSTRATE	2 = SEMI-ERECT 3 = ERECT				
16. SEED:					
1 Shape: l = OVATE	2 = OVAL 3 = ELLIPTICAL	Cheek: 1 = ROUNDED	2 = ANGULAR		
1 Brush: 1 ≈ SHORT	2 = MEDIUM 3 = LONG	1 Brush: 1 = NOT COL	LARED 2 = COLLARED		
Phenol reaction 1 = IVORY 2 = FAWN 3 = LT. BROWN (See instructions): 4 = BROWN 5 = BLACK					
3 Color: 1 = WHITE 2	2 ÷ AMBER 3 ≈ RED 4 = PURPLE	5 = OTHER (Specify)			
0 6 MM. LENGTH	0 4 MM. WIDTH	3 3 GM. PER 1000 SI	EEDS		
17. SEED CREASE:					
_ Width:] = 60% OR LE	SS OF KERNEL 'WINOKA'	Depth: 1 = 20% OR	LESS OF KERNEL 'SCOUT'		
1 1 1	S OF KERNEL 'CHRIS'		LESS OF KERNEL 'CHRIS'		
= :	WIDE AS KERNEL 'LEMHI'	3 = 50% OR I	LESS OF KERNEL 'LEMHI'		
	d, 1 = Susceptible, 2 = Resistant)	<u> </u>			
1 STEM RUST (Races) 15B-2,15]	LEAF RUST	STRIPE RUST	2 LOOSE SMUT		
151,32,31,113 POWDERY MILDEW	688, UN2-70-22, BUNT UN0168A	2 OTHER (Specify) S	pindle streak mosiac virus		
10 INSECT: (0 = Not Tosted	, 1 = Susceptible, 2 = Resistant)				
0 SAWFLY	0 APHID (Bydv.)	0 GREEN BUG	1 CEREAL LEAF BEETLE		
OTHER (Specify)	HESSIAN FLY	2 GP 2 A	1 B 2 c		
	RACES: (1 D 1 E	2 F 1 G		
20. INDICATE WHICH VARIETY MOST CLOSELY RESEMBLES THAT SUBMITTED:					
CHARACTER	NAME OF VARIETY	CHARACTER	NAME OF VARIETY		
Plant tillering	Lucas	Seed size	Lucas		
	Lucas	Seed shape	Lucas		
Leaf size	Lucas	Coleoptile elongation			
	Lucas		Lucas		
Leaf carriage	Lugan	Seedling pigmentation	Redcoat		

INSTRUCTIONS

GENERAL: The following publications may be used as a reference aid for the standardization of terms and procedures for completing this form:

- (a) L.W. Briggle and L. P. Reitz, 1963, Classification of Triticum Species and Wheat Varieties Grown in the United States, Technical Bulletin 1278, United States Department of Agriculture.
- (b) W.E. Walls, 1965, A Standardized Phenol Method for Testing Wheat Seeds for Varietal Purity, contribution No. 28 to the handbook of seed testing prepared by the Association of Official Seed Analysts. (See attachment.)

LEAF COLOR: Nickerson's or any recognized color fan should be used to determine the leaf color of the described variety.

FORM GR-470-6 (2-15-73)

10. LEAF:

3

2

2

Flag leaf at 1 = ERECT 2 = RECURVED booting stage: 3 = OTHER (Specify): Semi-erect

MM. LEAF WIDTH (First leaf below flag leaf)

2 = PRESENT

2

Hairs of first leaf sheath: 1 = ABSENT

UNITED STATES DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE GRAIN DIVISION HYATTSVILLE, MARYLAND 20782

1 Flag leaf: 1 = NOT TWISTED 2 = TWISTED

Waxy bloom of flag leaf sheath: 1 = ABSENT

7 CM. LEAF LENGTH (First leaf below flag leaf):

EXHIBIT C (Wheat)

OBJECTIVE DESCRIP	
INSTRUCTIONS: See Reverse.	FOR OFFICIAL USE ONLY
Ohio Agricultural Research & Development Co	B VPO NUMBER
ADDRESS (Street and No. or R.F.D. No., City, State, and ZIP Code)	740078 NHE
Wooster, Ohio 44691	VARIETY NAME OR TEMPORARY
WOOStel, Onlo 44091	DESIGNATION
·	Rulerate
Place the appropriate number that describes the varietal character	of this variety in the boxes below.
Place a zero in first box (e.s. 0 8 9 or 0 9) when number i	s either 99 or less or 9 or less.
1. KIND:	
1 = COMMON 2 = DURUM 3 = EMMER 4 = SPELT 5	= POLISH 6 ± POULARD 7 € CLUB
2. TYPE: 2 1 = SPRING 2 = WINTER 3 = OTHER (Specify)	1 = SOFT 3 = OTHER (Specify) 2 = HARD
2] = WHITE 2 = RED 3 = OTHER (Specify)	
3. SEASON - NUMBER OF DAYS FROM EMERGENCE TO:	
2 3 0 FIRST FLOWERING	2 3 5 LAST FLOWERING
4. MATURITY (50% Flowering):	
NO. OF DAYS EARLIER THAN	1 = ARTHUR 2 = SCOUT 3 = CHRIS
0 5 NO. OF DAYS LATER THAN	4 = LEMHI 5 = NUGAINES 6 = LEEDS
5. PLANT HEIGHT (From soil level to top of head):	
0 9 9 cm. HIGH	
0 4 CM. TALLER THAN	
CM. SHORTER THAN	4 = LEMHI 5 = NUGAINES 6 = LEEDS
6. PLANT COLOR AT BOOTING (See reverse):	7. ANTHER COLOR:
3 1 = YELLOW GREEN 2 = GREEN 3 = BLUE GREEN	1 1 = YELLOW 2 = PURPLE
8. STEM:	
Anthocyanin: 1 = ABSENT 2 = PRESENT	1 Waxy bloom: 1 = ABSENT 2 = PRESENT
Hairiness of last internode of rachis: 1 = ABSENT 2 = PRESENT	1 Internodes: 1 = HOLLOW 2 = SOLID
0 4 NO. OF NODES (Originating from node above ground)	2 2 CM. INTERNODE LENGTH BETWEEN FLAG LEAF
9. AURICLES:	
1 Anthocyanin: 1 = ABSENT 2 = PRESENT	Hairiness: 1 = ABSENT 2 = PRESENT